DEPARTMENT OF THE NAVY BUREAU OF NAVAL PERSONNEL WASHINGTON, DC 20370-5000

BUPERSINST 1000.22 Pers-23 12 February 1993

BUPERS INSTRUTION 1000.22

From: Chief of Naval Personnel

To: All Ships and Stations (less Marine Corps field addressees not having Navy personnel attached)

Subj: MANAGEMENT AND DISPOSITION OF NAVY PERSONNEL WITH PSEUDOFOLLICULITIS BARBAE (PFB)

Ref: (a) U. S. Navy Uniform Regulations

(b) OPNAVINST 5100. 19B (NOTAL)

Encl: (1) Pseudofolliculitis Barbae Treatment Protocol

(2) Waiver/Evaluation/Disposition

- 1. **Purpose**. To provide standardization in the management and disposition of service members with Pseudofolliculitis Barbae (PFB).
- 2. Background. Grooming standards to promote uniformity in the military services have included the requirement to be clean shaven. Reference (a) permits the wearing of mustaches and allows for facial hair for certain medical conditions such as uncontrollable PFB (shaving bumps), severe facial scarring, etc. Reference (b) describes the associated problems of facial hair which may constitute a safety or health hazard for some personnel.
- 3. **Discussion**. PFB is not a medical problem unless an individual is required to shave daily. This condition affects mainly the neck area but can affect any area of shaved skin. It occurs predominantly in blacks, but members of other races with strongly curved hair may also develop this condition. PFB is a foreign body type of skin reaction. As the sharp lip of closely shaved hair grows out, it curves sharply, forms a loop and penetrates the skin adjacent to the hair follicle. Shaving causes or aggravates the disorder, as only short stiff hairs reenter the skin. No conventional system or schedule of shaving can cure this disorder. However, in most cases it can be controlled.

- 4. **Policy**. Medical Department personnel must take necessary steps to educate both the patients and their supervisors regarding the nature and treatment of PFB. Supervisors of individual who are diagnosed with PFB must actively monitor and ensure treatment regimen is followed for those individuals as outlined in this instruction. It is the intent of this policy that a one-time thorough evaluation of each affected member's condition be sufficient. Members will not be required to be reevaluated at any tine or recommended for separation, subsequent to final determination, including transfer.
- Treatment. Individuals who state they cannot shave because of facial irritation will be referred to a medical officer for evaluation, diagnosis, and treatment. Enclosure (1) contains the treatment protocol that will be followed by all individuals with a diagnosis of PFB. After a PFB diagnosis has been made, the physician or Senior Medical Department Representative (SMDR) will instruct the patient and designated supervisor on the management and treatment protocol for PFB. Enclosure (2) will be utilized to document the service member's initial condition, progression of treatment, and final recommendation. The original will be placed in the member's health record and a copy will be placed in the member's service record. Individuals who have completed and failed the PFB treatment protocol will be recommended for a permanent "no shave" status. Enclosure (2) must be completed and presented to the commanding officer for action.

6. Action

- a. <u>Individuals</u> in a temporary or permanent "no shave" status must:
- (1) Maintain facial hair not to exceed $\frac{1}{4}$ inch in length. The beard will not become unkept in appearance and will not be manicured to any style.
 - (2) Comply with all treatment protocols.
- b. <u>Commands</u> will place emphasis on education, monitoring, and assistance when member is first treated. If the commanding officer determines that a permanent "no shaving" status is detrimental to good order and discipline or affects the member's ability to perform military duties, he/she <u>may</u> submit a <u>recommendation</u> to the Bureau of Naval Personnel (BUPERS) (Pers-282) for "Administrative

Separation By Reason of Pseudofolliculitis Barbae (PFB)."

In all cases, the final determination rests with BUPERS

(Pers-2). The recommendation will contain the following information:

- (1) A certified copy of the Shaving Waiver/Evaluation/Disposition Form (enclosure 2).
- (2) Commanding officer comments relative to his/her recommendation. Documentation of command impact of inability to perform military duties must be provided.
- (3) A statement from the service member, if so desired. If member declines to submit a statement, it must be so stated.
- (4) A "close-up" photograph of good quality which clearly depicts the member's PFB lesions.

c. BUPERS (Pers-282) will:

- (1) Consider each case on a "case by case" basis.
- (2) Notify the parent command of final disposition.
- d. <u>BUPERS (Pers-61)</u> will monitor the program as part of the Navy Affirmative Action Plan (NAAP).
- e. The Medical Officer and the member's command will document the final determination in the health and service records of the member. A copy of enclosure (2) will be filed in the service record, below the Performance Data Section in the case of enlisted members, and chronologically with service record page 13's, in the case of officers.
- 7. Form. NAVPERS 1000/1 (1-93), Pseudofolliculitis Barbae (PFB) Shaving Wavier/Evaluation/Disposition, will be available from BUPERS, Recruiting Programs and Policy Branch (PER-231), Washington, DC 20370-2310.

Blutop

BUPERSINST 1000.22 12 FEB 1993

Distribution: SNDL Parts 1 and 2

Stocked:

Naval Aviation Supply Office Physical Distribution Division Code 103 5801 Tabor Avenue Philadelphia PA 19120-4099 (500 copies)

PSEUDOFOLLICULITIS BARBAE (PFB) TREATMENT PROTOCOL

PHASE I: Control of mild cases (less than 20-30 small papules or rare pustules)

- 1. Totally avoid shaving for 3 to 4 weeks until all lesions have subsided, while applying Vioform-HC cream (NSN 6505-00-074-9912) to involved skin each morning.
- 2. To soften whiskers, begin a nightly application of Retin-A Cream 0.05 0.1% (NSN 6505-01-044-9389, .05%) (NSN 6505-01-044-9388, .1%) to beard skin while beard is growing out. Use as tolerated.
- 3. Circular brushing of beard area with soft brush (polyester skin cleaning pad) or medium firm toothbrush, four times a day for 3 to 5 minutes.
- 4. Once shaving bumps subside, avoid close shave by following the steps below.
- a. Water soften beard first with hot, wet washcloth for 5 minutes.
 - b. Use lubricating shaving gel.
 - c. Avoid close shave by using a PFB Razor.
- d. Shave with the grain of the beard and do not stretch the skin. Use only one stroke over each area of the beard.
 - e. Shave only every 2 to 3 days.
- 5. Instead of following the steps in paragraph (4), use an electric razor on high setting to avoid close shaving and:
 - a. Prepare beard with electric razor pre-shave.
 - b. Do not stretch skin while shaving.
 - c. Shave with the grain of beard growth.
- d. Avoid multiple repetition of strokes in the same area.
 - e. Do not Dress razor head hard against the skin.

- f. Shave every other day.
- 6. After shaving, rub in Vioform-HC cream at the first sign of irritation, tenderness or new papule formation.

PHASE II: Control of moderate cases (20-60 papules with pustules) or cases unresponsive to Phase I

- 1. Continue Phase I steps 1, 2, and 3.
- 2. To remove whisker and leave blunted end, use either (a) or (b) below:
 - a. Barium sulfide depilatory very mild to strong.
 - (1) Use only every 48-72 hours.
 - (2) Use mildest form that works.
- (3) Apply to small areas at a time (1/4 1/2) of face) and start removal no later than 3 minutes after application.
- (4) Remove with spatula, butter knife, tongue blade, or other blunt straight edge.
- (5) Use short, rapid strokes against the direction of hair growth. Keep shaving instrument moist but wiped clean.
- (6) If some hairs remain, repeat after 30-60 seconds. Wait if within 3 minutes of original application.
- (7) Rapid and thorough rinsing, with soap washes between rinses, two to three times daily
- (8) Neutralize depilatory with a diluted vinegar solution (1 teaspoon per pint of water) then rinse again thoroughly with tap water.
- (9) Apply Vioform-HC cream if irritation develops.
 - b. Calcium triglycolate depilatory.
 - (1) Apply to bearded areas.
 - (2) Leave on 5 to 10 minutes.
- (3) Use spatula, tongue blade, butter knife, or other blunt straight edge to remove material with strokes against the direction of hair growth.

- (4) Rinse thoroughly 2-3 times.
- $\mbox{(5)}$ Apply Vioform-HC cream if irritation develops.

PHASE III: Control of severe cases or in those cases unresponsive to Phases I and II

- 1. Continue Phase I, steps 1, 2, 3, and 6.
- 2. Use barber's clipper (with clipping guard) to prevent a closer than 1mm cut of whisker hair.
- 3. Hold clipper close to face and slowly glide along facial contours against the direction of hair growth.
- 4. Do not stretch skin nor press hard against the skin with the clippers.
- 5. Use clippers on a daily basis.

PSEUDOFOLLICULITIS BARBAE (PFB) SHAVING WAIVER/EVALUATION/DISPOSITION

NAI	ME	(LAST)	(FIRST)	Μ.	I.		SS	SN	
ME1	DICAL	OFFICER INI	TIAL EVALU	ATIO	<u>N</u>				
((() Ps	PFB or othe eudofollicul cial Noduloc her:	itis Barba ystic Acne	e (Pi	FB)			shavi	ng
NAI	ME OF	MEDICAL OFF	ICER/RANK			SIG	NATURE/DA	TE	
		the medical hair is reco							G of
SPECIFY PERIOD OF TIME TITLE/SIGNATURE/DATE									
MEI	DICAL	OFFICER//SM	DR DOCUMEN	TATI(ON OF	PFB I	PROTOCOL	COMPLE	TION
() PH	ASE I				_			
			SIGNATURE	/TIT	LE		DA	ΛTE	
() PH	ASE II							
			SIGNATURE	/TIL	E		DA	TE	
() PH	ASE III							
`	,		SIGNATURE	/TIT	LE		DA	TE	
FA:	ILURE	OF PFB PROT	OCOL RECOMI	MEND	ATION				
		vy member ha nt "No Shave					PFB prot	ocol.	A
		SIGN	ATURE/ TIT	LE			DA	TE	
COI	MMAND	ING OFFICER	DECISION:						
(permanent "N fer to BUPER							
		NAME/ RAN	K/ TITLE			SIC	GNATURE/	DATE	